

Approximation of unbounded functions by trigonometric polynomials

ALAA A. AUAD

alaa.adnan66.aa@gmail.com

University Of Al-Anbar ,
Faculty Of Education

MOUSA M. KHRAJAN

mmkrady@gmail.com

University Of Al-Muthana,
Science College

Abstract

The purpose of this paper is present a brief survey of know on estimates the rate for best approximation of unbounded functions by suitable trigonometric polynomials of one variable in weighted space $L_{p,\alpha}(X)$. Moreover two theorems (3) and (4) concerning the rate of trigonometric approximation of $f^{(k)}$ with k non-integer in $L_{p,\alpha}(X)$.

Keywords: Unbounded functions, trigonometric polynomials, weight space

تقريب الدوال الغير مقيدة بواسطة متعددات الحدود المثلثية

: موسى مكي خريجان

جامعة المثنى – كلية العلوم

mmkrady@gmail.com

الباحث : علاء عدنان عواد الباحث

جامعة الانبار – كلية التربية للعلوم الصرفة

alaa.adnan66.aa@gmail.com

المستخلص

الغرض من هذا البحث هو عرض دراسة موجزة لمعرفة تخمين درجة افضل تقريب للدوال الغير مقيدة في الفضاء الموزون بواسطة متعددات الحدود المثلثية ذات المتغير الواحد. اضافة الى ذلك المبرهنات الثالثة والرابعة تتعلق بالتقريب المثلثي للدوال $f^{(k)}$ في فضاء الوزون حيث ان k عدد غير صحيح.

1. Introduction

In 1977 Taberski R. see [10] and in 1979 Andreev, Popov, Sendov see [2] introduced the best approximation of functions by trigonometric polynomials of fractional order. This notion could be considered as direct generalization of classical trigonometric approximation in weighted space $L_{p,\alpha}(X)$ and it is more natural to use it for number of problems of approximation functions (see, for example [3],[5],[6], [9]).

The important problem of approximation theory and theory of Fourier series is the problem of description of best approximation of functions by trigonometric polynomials see [7] and [10].

One can consider this problem from the viewpoint of description of Roman Taberski of trigonometric approximation. In this paper, we obtain the description of some estimates for the best trigonometric approximation of unbounded functions in $L_{p,\alpha}(X)$ and the order approximation of $f^{(k)}$ by suitable trigonometric polynomials.

Let $X = [-\pi, \pi]$, we denote the $L_p(X)$ -norm ($1 \leq p < \infty$),

$$\|f\|_p = \left(\int |f(x)|^p dx \right)^{\frac{1}{p}} < \infty \dots\dots\dots(1)$$

and define for suitable $W(\alpha, x)$ of all weight function on closed interval X such that $|f(x)| \leq M\alpha(x)$, where M is positive real number and $\alpha: X \rightarrow \mathbb{R}^+$ weight function.

We shall denote $L_{p,\alpha}(X)$ the space of all unbounded functions on X which are equipped with the following norm

$$\|f\|_{p,\alpha} = \left(\int_X \left| \frac{f(x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} < \infty \dots\dots\dots(2).$$

Let H_n be the set of all 2π – periodic trigonometric polynomials of order less than or equal n , $n \in \mathbb{N} = \{1, 2, 3, \dots\}$.

The degree of best trigonometric approximation of an arbitrary function in weighted space $L_{p,\alpha}(X)$ is define by

$$E_n(f)_{p,\alpha} = \inf \{ \|f - t_n\|_{p,\alpha} , \quad t_n \in H_n \} \dots\dots\dots(3).$$

Denote by $\omega(f, \delta)_{p,\alpha}$ the modulus of continuity with respect to the $L_{p,\alpha}$ –norm i.e.

$$\omega(f, \delta)_{p,\alpha} = \sup_{h \in [0, \delta]} \|\Delta_h^1 f\|_{p,\alpha} \quad (0 \leq \delta < \infty) \dots\dots\dots(4)$$

where $\Delta_h^1 f(x) = |f(x) - f(x + h)|$.

$$\text{Suppose that } F(f) = \sum_{m=-\infty}^{\infty} C_m \frac{e^{imx}}{\alpha(x)} \dots\dots\dots(5)$$

is the Fourier series of $f \in L_{p,\alpha}(X)$ for which the integral over X is zero so that $C_0 = 0$. Given any $a > 0$, we define the a -th integers of f by the identity

$$I_a(f, x) = \sum_{m=-\infty}^{\infty} \frac{C_m}{(im)^a} \frac{e^{imx}}{\alpha(x)} \dots\dots\dots(6),$$

where $\frac{1}{(im)^a} = |m|^{-a} \exp(-\frac{1}{2}\pi i a \text{ sign } m)$.

As is well known [12] for $f_a(x) = I_a(f, x)$ exist possibly for almost every x , is Lebesgue-integrable and $F[f_a] = f_a(x)$ a.e.

In this case for $a > 0$, the convolution

$$f_a(x) = (f * I_a) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{f(x-u)I_a(u)}{\alpha(x)} du \dots\dots\dots(7)$$

is of class $L_{p,\alpha}(X)$.

In several cases, the last Fourier series converges for every or a.e. x , its sum

$$f^{(-a)}(x) \equiv I_a[f](x) \text{ see [12].}$$

If $0 < a < 1$, the derivative $f(x)$ of f is defined by the formula

$$f^{(a)}(x) = \frac{d}{dx} I_{1-a}(f, x) \dots\dots\dots(7)$$

provided the right said exists. We set

$$f^{(a+k)}(x) = (f^{(a)}(x))^k = \frac{d^{k+1}}{dx^{k+1}} I_{1-a}(f, x) \dots\dots\dots(8)$$

For positive integer k .

Let f be a complex-valued function defined in the closed interval $[-\pi, \pi]$.
Then

$$V_{p,\alpha}(f, -\pi, \pi) = \sup_{\Pi} \left\{ \sum_{j=0}^{n-1} |f(x_{j+1}) - f(x_j)|^p \right\}^{\frac{1}{p}} < \infty, (1 \leq p < \infty) \dots\dots\dots(9)$$

where Π denotes the partition $\{-\pi = x_0 < x_1 < \dots < x_n = \pi\}$ is often called the p -th variation of f in $[-\pi, \pi]$.

The aim of this paper is to present some approximation theorems for unbounded functions in space $L_{p,\alpha}(X)$. Their proofs are based on the suitable result announced in [4] and [11].

2. Auxiliary theorem

Let us explicitly formulate direct theorem of the order degree of best approximation of unbounded functions by algebraic or trigonometric polynomials in weighed space.

Theorem (A):[1]

Let f be unbounded function in weighted space $L_{p,\alpha}(X)$ and $(1 \leq p < \infty)$. For every natural k there exist constant $c(k)$ depending on k , such that

$$E_n(f)_{p,\alpha} \leq c(k) \omega_k(f, \sqrt{n})_{p,\alpha}.$$

3. Main results

Considering for any unbounded function f belong to weighted space $L_{p,\alpha}(X)$ such that $\frac{f(x)}{\alpha(x)} \leq M$ for all $x \in X$, we shall prove the following

Theorem (1)

Suppose that function f in space $L_{p,\alpha}(X)$, $(1 \leq p < \infty)$, with a derivative $f^{(k-1)}$ of non-negative integer order $k - 1$ and absolutely continuous in X . Then

$$E_n(f)_{p,\alpha} \leq \frac{C(k)}{n^k} E_n(f^k)_{p,\alpha} \text{ for } n = 0, 1, 2, \dots$$

Proof :

Let (5) be the Fourier series of $f \in L_{p,\alpha}(X)$ and k positive integer.

We have

$$f^{(k)}(x) = \frac{d^k}{dx^k} I_k[f](x) = \sum_{m=-\infty}^{\infty} C_m(im)^k \frac{e^{imx}}{\alpha^{(k)}(x)} \text{ for all } x. \text{ Moreover}$$

$$F[f^{(k)}](x) = \sum_{m=-\infty}^{\infty} C_m(im)^k \frac{e^{imx}}{\alpha^{(k)}(x)}.$$

$$\text{Thus } f^{(k)}(x) = \sum_{m=-\infty}^{\infty} C_m(im)^k \frac{e^{imx}}{\alpha^{(k)}(x)} = F[f^{(k)} * \Phi_n](x) \text{ where}$$

$$\Phi_n(x) = \sum_{m=-\infty}^{\infty} \frac{e^{imx}}{(im)^k} = 2 \sum_{m=1}^{\infty} \frac{\sin mx}{m}.$$

For each trigonometric polynomials $\xi_n(x), \eta_n(x)$ of order n at most, there is a trigonometric polynomial $T_n(x)$ such that

$$\begin{aligned} f(x) - T_n(x) &= \frac{1}{2\pi} \int_X \{f^{(k)}(t) - \xi_n(t)\} \{\Phi_n(x-t) - \eta_n(x-t)\} dt \\ &= \frac{1}{2\pi} \int_X \{f^{(k)}(x-u) - \xi_n(x-u)\} \{\Phi_n(u) - \eta_n(u)\} du. \end{aligned}$$

Thus

$$E_n(f)_{p,\alpha} \leq \|f - T_n\|_{p,\alpha} = \left(\int_X \left| \frac{f(x) - T_n(x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}}$$

$$\leq \frac{1}{2\pi} \left\{ \left(\int_X \left| \frac{f^{(k)}(x-u) - \xi_n(x-u)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} \cdot \left(\int_X \left| \frac{\Phi_n(u) - \eta_n(u)}{\alpha(u)} \right| du \right) \right\}$$

whence

$$E_n(f)_{p,\alpha} \leq \frac{1}{2\pi} E_n(f)_{p,\alpha} \cdot E_n(\Phi_n)_{1,\alpha}.$$

We have from direct theorem (A)

$$E_n(f)_{p,\alpha} \leq C \omega(f, \frac{1}{n+1})_{1,\alpha},$$

from properties of modulus of continuity, we have

$$\omega(f, \lambda \delta) = \lambda \omega(f, \delta),$$

$$\text{so, } \Phi_n(u) = \begin{cases} \pi - u & \text{if } 0 < u < \pi \\ 0 & \text{if } u = 0, u = \pi \end{cases}$$

$$\text{i.e. } V(\Phi)(u) = 2(2\pi) = 4\pi, \quad u \in [-\pi, \pi].$$

Hence

$$E_n(f)_{p,\alpha} \leq \frac{4\pi C_1(p)}{n+1} \cdot \frac{1}{2\pi} E_n(f^{(k)})_{p,\alpha} \leq \frac{C_2(p)}{n+1} \cdot E_n(f^{(k)})_{p,\alpha}.$$

Theorem (2)

Let $f \in L_{p,\alpha}(X)$, $(1 \leq p < \infty)$ and the trigonometric polynomial $T_n(x) = T_n(f, x)$ of order n at most such that

$$\|f - T_n\|_{p,\alpha} \leq C(p) E_n(f)_{p,\alpha}, \quad n = 0, 1, 2, \dots \quad \dots\dots\dots(10)$$

for all satisfies the conditions of theorem (1), with positive integer k . Then

$$\|f^{(k)} - T_n^{(k)}(f)\|_{p,\alpha} \leq C(p, k) E_n(f^{(k)})_{p,\alpha}, \quad n = 0, 1, 2, \dots$$

.....(11).

Proof :

Let $F_v(f, x)$ be denote the $v - th$ partial sum of the Fourier series of f and let $V_n(f, x)$ be the Vallee-poussin means of this series, defined by the foemula

$$V_n(f, x) = \frac{1}{n+1} \sum_{v=n}^{2n} F_v(f, x), \quad n = 0, 1, 2, \dots$$

As is well know, for any $f \in L_{p,\alpha}(X)$

$$V_n(f, x) = \frac{1}{\pi} \int_X \left| \frac{f(x+t)\beta_n(t)}{\alpha(t)} \right| dt, \quad \text{if } p = 1$$

$$\text{where } \beta_n(t) = \frac{1}{\pi} \sum_{v=n}^{2n} \frac{\sin(v+\frac{1}{2})t}{2\sin\frac{1}{2}t} = \frac{\sin(n+\frac{1}{2})t \cdot \sin(\frac{3}{2}n+\frac{1}{2})t}{2(n+1)(\sin\frac{1}{2}t)^2} \text{ and}$$

$$\frac{1}{\pi} \int_X \beta_n(t) dt = 1, \frac{1}{\pi} \int_X |\beta_n(t)| dt < 2 \cdot \frac{4n+1}{2n+2} < 4.$$

Moreover

$$\psi_n(f^{(k)}, x) = \psi_n^{(k)}(f, x), \text{ whenever } f^{(k)} \in L_{p,\alpha}(X).$$

By Minkowsk's inequality, we obtain

$$\begin{aligned} \|f^{(k)} - T_n^{(k)}(f)\|_{p,\alpha} &= \left(\int_X \left| \frac{f^{(k)}(x) - T_n^{(k)}(f, x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} \\ &\leq \left(\int_X \left| \frac{f^{(k)}(x) - \psi_n^{(k)}(f, x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} + \\ &\quad \left(\int_X \left| \frac{\psi_n^{(k)}(f, x) - T_n^{(k)}(\psi_n(f, x))}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} \end{aligned}$$

$$+ \left(\int_X \left| \frac{T_n^{(k)}(\psi_n(f,x)) - T_n^{(k)}(f,x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}}$$

Set $p_n(f, x)$ the trigonometric polynomial of best approximation of f in weighted space $L_{p,\alpha}(X)$ of order n , we have

$$\begin{aligned} & \left(\int_X \left| \frac{f^{(k)}(x) - \psi_n^{(k)}(f,x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} \leq \\ & \left(\int_X \left| \frac{f^{(k)}(x) - p_n^{(k)}(f,x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} + \left(\int_X \left| \frac{p_n^{(k)}(f,x) - \psi_n^{(k)}(f,x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} \\ & \leq \|f^{(k)} - p_n(f^{(k)})\|_{p,\alpha} + \|\psi_n(p_n(f^{(k)})) - f^{(k)}\|_{p,\alpha} \\ & \leq C_1 E_n(f^{(k)})_{p,\alpha} \dots\dots\dots(12) \end{aligned}$$

from Zygmund inequality gives

$$\begin{aligned} & \left(\int_X \left| \frac{T_n^{(k)}(\psi_n(f,x)) - T_n^{(k)}(f,x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} \leq \\ & n^k \left(\int_X \left| \frac{T_n(\psi_n(f,x)) - T_n(f,x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} \dots\dots\dots(13) \end{aligned}$$

And

$$\begin{aligned} & \left(\int_X \left| \frac{\psi_n^{(k)}(f,x) - T_n^{(k)}(\psi_n(f,x))}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} \leq (2n)^k \left(\int_X \left| \frac{\psi_n(f,x) - T_n(\psi_n(f,x))}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} \\ & \leq C_2(p)(2n)^k E_n(\psi_n(f))_{p,\alpha} \dots\dots\dots(14) \end{aligned}$$

From (13) and (14), we obtain

$$\left(\int_X \left| \frac{T_n(\psi_n(f,x)) - T_n(f,x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} \leq \left(\int_X \left| \frac{T_n(\psi_n(f,x)) - \psi_n(f,x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}}$$

$$+ \left(\int_X \left| \frac{\psi_n(f, x) - f(x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}} +$$

$$\left(\int_X \left| \frac{f(x) - T_n(f, x)}{\alpha(x)} \right|^p dx \right)^{\frac{1}{p}}$$

$$\leq C_3(p) E_n(\psi_n(f))_{p, \alpha} + C_4 E_n(f)_{p, \alpha} + C_5(p) E_n(f)_{p, \alpha} \dots (15)$$

and

$$E_n(\psi_n(f))_{p, \alpha} \leq C E_n(f)_{p, \alpha}$$

.....(16).

Thus, from (12), (15) and (16), we get

$$\|f^{(k)} - T^{(k)}\|_{p, \alpha} \leq C_6 E_n(f^{(k)})_{p, \alpha} + C_7 n^{(k)} (C_8(p) + 1) E_n(f)_{p, \alpha}$$

$$+ C_9(p) E_n(f)_{p, \alpha}$$

and by theorem (1) then (11) follows.

Theorem (3)

Let $f \in L_{p, \alpha}(X)$, $(1 \leq p < \infty)$ and that the derivative $f^{(a)}$ in weighed space with $0 < a < 1$. Then

$$E_n(f)_{p, \alpha} \leq \frac{C(a)}{(n+1)^a} E_n(f^{(a)})_{p, \alpha} \text{ for } n = 0, 1, 2 \dots \dots \dots (17).$$

Proof :

$$\text{Putting } (x) = \int_0^x f(u) du, \text{ we have } G_n(x) = \sum_{m=-\infty}^{\infty} C_m(im)^{-1} (e^{imx})$$

$$= f_1(x) - f(0)$$

uniformly in $x \in -\infty, \infty$. Consequently $G_n(x) = I_n(x, g) - I_n(0, g) \quad \forall x$.

Therefore, for all real x

$$G_n(x) = \frac{1}{\pi} \int_X \Phi_n(x-u)g(u)du - f(0),$$

$$\int_X \Phi_n(x-u)du = 0.$$

For each trigonometric polynomials $\zeta_n(x)$ and $\eta_n(x)$ of order n , there is a trigonometric polynomial $T_n(x)$ such that

$$G_n(x) - T_n(x) = \frac{1}{2\pi} \int_X \{g(x-u) - \zeta_n(x-u)\} \cdot \{\Phi_a(u) - \eta_n(u)\}du - I_a(0, g).$$

Assuming $0 < b < \pi$, we have

$$\begin{aligned} & \left\{ \int \left| \frac{G(x+b)-T_n(x+b)-(G(x-b)-T_n(x-b))}{2b \alpha(x)} \right|^p dx \right\}^{\frac{1}{p}} = \\ & \frac{1}{2\pi} \left\{ \int_X \left| \int_X \frac{g(x+b-u)-\zeta_n(x+b-u)-(g(x-b-u)-\zeta_n(x-b-u))}{2b \alpha(x)} \cdot \{\Phi_a(u) - \right. \right. \\ & \left. \left. \eta_n(u)\} du \right|^p dx \right\}^{\frac{1}{p}} \\ & \leq \frac{1}{2\pi} \int_X |\Phi_a(u) - \eta_n(u)| \cdot \left\{ \int_X \left| \frac{g(x+b-u)-\zeta_n(x+b-u)-(g(x-b-u)-\zeta_n(x-b-u))}{2b \alpha(x)} \right|^p dx \right\}^{\frac{1}{p}} du \end{aligned}$$

we take the best approximation of functions $g(v)$ and $\Phi_a(v)$ they $\zeta_n(u)$, $\eta_n(u)$ in spaces $L_{p,\alpha}(X)$ and $L_1(X)$ respectively. Then

$$\begin{aligned} & 2\pi \left\{ \int_X \left| \frac{G(x+b)-G(x-b)}{2b \alpha(x)} - \frac{T_n(x+b)-T_n(x-b)}{2b \alpha(x)} \right|^p dx \right\}^{\frac{1}{p}} \\ & \leq \left\{ \int_X \left| \frac{g(v+b)-g(v-b)}{2b \alpha(x)} - \frac{\zeta_n(v+b)-\zeta_n(v-b)}{2b \alpha(x)} \right|^p dv \right\}^{\frac{1}{p}} \cdot E_n(\Phi_a)_1. \end{aligned}$$

By theorem (2), we have

$$2\pi \left\{ \int_X \left| \frac{G_n^{(1)}(x) - T_n^{(1)}(x)}{\alpha(x)} \right|^p dx \right\}^{\frac{1}{p}} \leq \left\{ \int_X \left| \frac{g^{(1)}(v) - \zeta_n^{(1)}(v)}{\alpha(v)} \right|^p dv \right\}^{\frac{1}{p}} \cdot E_n(\Phi_a)_1 .$$

$$\leq C(p) E_n(g^{(1)})_{p,\alpha} \cdot E_n(\Phi_a)_1$$

So, $\|f - T_n^{(1)}\|_{p,\alpha} \leq 2\pi \left\{ \int_X \left| \frac{f(x) - T_n^{(1)}(x)}{\alpha(x)} \right|^p dx \right\}^{\frac{1}{p}}$ and consequently

$$E_n(f)_{p,\alpha} \leq \frac{1}{2\pi} C E_n(f^{(a)})_{p,\alpha} \cdot E_n(\Phi_a)_1 ,$$

By direct theorem [], we have

$$E_n(f)_{p,\alpha} \leq \frac{3}{\pi} C E_n(f^{(a)})_{p,\alpha} \omega_1\left(\frac{1}{n+1}, \Phi_a\right) \leq \frac{3}{\pi} C E_n(f^{(a)})_{p,\alpha} V_1(\Phi_a) \cdot \frac{1}{n+1}$$

$$\leq \frac{C(a)}{(n+1)^a} E_n(f^{(a)})_{p,\alpha} ,$$

the desired assertion is established.

Theorem (4)

Given any $f \in L_{p,\alpha}(X)$, $(1 \leq p < \infty)$, let us consider the trigonometric polynomial $T_n(x) = T_n(f, x)$ of order n such that

$$\|f^{(k)} - T_n^{(k)}\|_{p,\alpha} \leq C(p, k) E_n(f^{(k)})_{p,\alpha} , \text{ with some positive non-integer } k.$$

The proof runs analogously of theorem (2).

References

1. A. Alaa, On the degree of best approximation of unbounded functions algebraic polynomials, Journal Ibn-Al-Haitham journal for pure and applied science, vol. 29, Iss. 2 (2016).
2. A. Andreev, V. Popov, B. Sendov, Jackson-type theorem for best one sided approximation by trigonometric polynomials and spline, Zameki, Vol. 26, No. 5 (1979), pp. 889-896.
3. A. Banbenko, N. Chennykh, V. Shevaldin, The Jackson-Stechkin inequality in L_2 with a trigonometric modulus of continuity, Math. Zameki, pp. 928-932 (in Russian)
English translation : Math. Notes 65 (5-6) (1999), pp. 777-781.
4. R. DeVore, G. Lorentz, Constructive approximation, Springer Verlag, Berlin (1993).
5. B. Deranganov, A. Parvanov, On estimating the rate of best trigonometric approximation by a modulus of smoothness, Acta. Math. Hungar, 131 (2011), pp. 360-379.
6. B. Deranganov, A new modulus of smoothness for trigonometric polynomial approximation, East journal approximation, 8 (4) (2002), pp. 465-499.
7. H. Meikle, A new twist to Fourier transform, Berlin, Dover (2004).
8. V. Popov, On convergence and saturation of Jackson polynomials in L_p - space, Journal approximation theory and application, 1-10 (1984).
9. V. Shevaldin, The Jackson-Stechkin inequality in the space $C(T)$ with trigonometric continuity modulus annihilating the first harmonics, proc. Steklov Inst. Math. Approximation theory, Asymptotic expansion, Suppl. 1 (2001), pp. 206-213.
10. R. Taberski, Approximation of functions possessing derivative of positive orders, Ann. Polon. Math. XXXIV (1977), pp. 13-23.
11. R. Taberski, Trigonometric approximation in the norms and semi norm, Stud. Math. T. LXXX (1984), pp. 197-217.
12. A. Zygmund, Trigonometric series, II, Cambridge (1959).